

Helps Prevent and Treat **Flu.**

**Flumivir®**  
Oseltamivir



Oseltamivir is FDA-approved for the treatment and prevention of adults with uncomplicated acute illness caused by influenza A or B viruses<sup>1</sup>.

**CDC – ACIP**

Recommends the use of Oseltamivir for Patients with Confirmed or suspected influenza <sup>1</sup>

Simple dosage and schedule with Flumivir® (Oseltamivir 75mg)<sup>1</sup>:

Treatment: 1 Capsule Twice Daily

Prevention: 1 Capsule Once Daily

Treatment BID		
	Dosage	Period
Adults (≥13 Years)	75 mg	5 days

Prevention OD		
	Dosage	Period
Adults (≥13 Years)	75 mg	7-10 days

Children Weight	Dosage	Period
≤15kg	30 mg	5 days
>15kg - 23kg	45 mg	5 days
>23kg - 40kg	60 mg	5 days
>40kg	75 mg	5 days

Children Weight	Dosage	Period
≤15kg	30 mg	7-10 days
>15kg - 23kg	45 mg	7-10 days
>23kg - 40kg	60 mg	7-10 days
>40kg	75 mg	7-10 days

References:

1. Antiviral Agents for the treatment and prophylaxis of Influenza, Recommendations of the advisory Committee on immunization practices (ACIP): CDC, MMWR/January 21,2011/Vol.60/No.1

Before prescribing please consult physicians' circular.

® Registered trade name



**CE GMP**



Benta Pharma Industries